

DFPD/93/TH/72

hep-th/9402081

NONPERTURBATIVE MODEL OF LIOUVILLE GRAVITY[¶]MARCO MATONE^{*}*Department of Physics “G. Galilei” - Istituto Nazionale di Fisica Nucleare**University of Padova**Via Marzolo, 8 - 35131 Padova, Italy***ABSTRACT**

We obtain nonperturbative results in the framework of continuous Liouville theory. In particular, we express the specific heat $\mathcal{Z}$ of pure gravity in terms of an expansion of integrals on moduli spaces of punctured Riemann spheres. The integrands are written in terms of the Liouville action. We show that $\mathcal{Z}$ satisfies the Painlevé I.

[¶]Partly supported by the European Community Research Programme *Gauge Theories, applied supersymmetry and quantum gravity*, contract SC1-CT92-0789

^{*}e-mail: matone@padova.infn.it, vaxfpd::matone

1. In this paper we introduce models of Liouville theory in the continuum which are based on the Riemann sphere with punctures. The models include pure gravity. In particular we will show that

$$\mathcal{Z}(t) = t^{-12} \sum_{k=4}^{\infty} t^{5k} \int_{\overline{\mathcal{M}}_{0,k}} \left(i\bar{\partial}\partial S_{cl}^{(k)} \right)^{k-4} \wedge \omega^{F_0} - \frac{t^3}{2} \quad (1)$$

is the specific heat of pure gravity, namely $\mathcal{Z}$ satisfies the Painlevé I

$$\mathcal{Z}^2(t) - \frac{1}{3}\mathcal{Z}''(t) = t. \quad (2)$$

$S_{cl}^{(k)}$ in (1) denotes the *classical* Liouville action on the k -punctured Riemann sphere. The class $[\omega^{F_0}]$ is the Poincaré dual of a divisor on the compactified moduli space $\overline{\mathcal{M}}_{0,k}$ which is given in terms of the $(2k-8)$ -cycles defining the Deligne-Knudsen-Mumford boundary of $\overline{\mathcal{M}}_{0,k}$. The basic tools to obtain (1) are classical Liouville theory and intersection theory.

This result reproduces in the continuum the well-known result obtained in the matrix model approach to pure gravity [1]. For reviews on matrix models and 2D gravity see [2].

2. The problems arising in the continuum formulation of Liouville gravity [3–5] are essentially:

- a. To evaluate Liouville correlators on Riemann surfaces of genus $h \geq 2$;
- b. To perform the integration on moduli spaces;
- c. To recover nonperturbative results from the topological expansion.

Results from matrix models and topological gravity show that these aspects are strictly related with the structure of $\mathcal{M}_h \equiv \mathcal{M}_{h,0}$ (we denote by $\mathcal{M}_{h,n}$ the moduli spaces of Riemann surfaces of genus h and n punctures). In particular, it turns out that the Liouville action is the Kähler potential for the natural (Weil-Petersson) metric on the moduli space. Also CFT is strictly related with the geometry of moduli space. For example, the Mumford isomorphism

$$\lambda_n \cong \lambda_1^{c_n}, \quad c_n = 6n^2 - 6n + 1,$$

where $\lambda_n = \det \text{ind } \bar{\partial}_n$ are the determinant line bundles, connects geometrical properties of $\mathcal{M}_h$ with the central charge $d = -2c_n$ of a weight n , b - c system (notice that $d \leq 1$). Actually, the bosonization of b - c systems can be used to reproduce the Coulomb gas formulation of $d \leq 1$ conformal matter. For $d > 1$ it is not possible to represent conformal matter by a b - c system. In this case one can consider the β - γ system of weight n whose central charge

is $2c_n$. However, the representation of the β - γ system in terms of free fields is a long-standing problem which seems related to the $d = 1$ barrier. These aspects indicate that there is a connection between the barrier and the Mumford isomorphism. This is related to a similar structure considered in [6] in the framework of the geometrical formulation of 2D gravity [6, 7] where representing elliptic and parabolic Liouville operators by means of a scalar field constrains the conformal matter to be in the sector $d \leq 1$.

The natural framework to investigate the aspects considered above is the theory of uniformization of Riemann surfaces where Liouville theory plays a crucial role. Actually, in [8] it has been shown that the Liouville action appears in the correlators (intersection numbers) of topological gravity [9]. The relationships between Liouville theory, matrix models and topological gravity suggest that it is possible to extend the above Liouville-topological gravity relationship by recovering the nonperturbative results of matrix models by continuum Liouville theory. In our model we will reduce all aspects concerning higher genus contributions to punctured spheres. The reduction to punctured sphere has been considered also by V.G. Knizhnik who expressed the sum of the genus expansion as a CFT on an arbitrary N -sheet covering of the Riemann sphere with branch points. For each branch point he associated a vertex operator and proposed to express the infinite sum on all genus ($h \geq 2$) as the limit for $N \rightarrow \infty$ of a ‘nonperturbative’ partition function [10].

A natural way to get punctured spheres is by pinching all handles of a compact Riemann surface. Degenerate (singular) surfaces belong to the boundary of moduli spaces. These singularities play a fundamental role in the evaluation of relevant integrals (intersection theory). The fact that the *classical* Liouville action is the Kähler potential for the Weil-Petersson metric and the structure of the boundary of moduli space suggest to consider integrals on $\overline{\mathcal{M}}_h$ in the framework of the Duistermaat-Heckman integration formula [11]. The final result should be a sum of integrals Z_n^F on the moduli space of punctured Riemann spheres $\overline{\mathcal{M}}_{0,n} = (\hat{\mathbf{C}} \setminus \Delta_n) / \text{Sym}(n) \times PSL(2, \mathbf{C})$ with the integrands involving the Liouville action. These remarks indicate that a theory à la Friedan-Shenker [12] can be concretely formulated to recover nonperturbative results in the continuum formulation. In this paper, we do not consider points **a-c** separately, rather we state the final solution finding the explicit form of the integrals Z_n^F on $\overline{\mathcal{M}}_{0,n}$.

The reduction to punctured sphere is particularly evident in topological field theory coupled to 2D gravity where higher genus contributions to the free energy $\langle 1 \rangle_h$ can be written in terms of the sphere amplitudes of the puncture operator P [9, 13]. The physical

observables of the theory are the primary fields $\mathcal{O}_\alpha$ ($\alpha = 0, 1, \dots, N-1$, $\mathcal{O}_0$ is the identity operator) and their gravitational descendents $\sigma_n(\mathcal{O}_\alpha)$, $n = 1, 2, \dots$. In the coupled system $\mathcal{O}_0$ becomes non-trivial and it is identified with P . Denoting by $\mathcal{L}_0$ the minimal Lagrangian, the more general one is $\mathcal{L} = \mathcal{L}_0 + \sum_{n,\alpha} t_{n,\alpha} \sigma_n(\mathcal{O}_\alpha)$, $\sigma_0(\mathcal{O}_\alpha) \equiv \mathcal{O}_\alpha$, where $t_{n,\alpha}$ are coupling constants. With this definition one can compute correlation functions with an insertion of σ_k just by differentiating $\langle 1 \rangle_h$ with respect to t_k . Thus in general

$$\langle \sigma_{d_1}(\mathcal{O}_{\alpha_1}) \cdots \sigma_{d_n}(\mathcal{O}_{\alpha_n}) \rangle_h = \frac{\partial}{\partial t_{d_1, \alpha_1}} \cdots \frac{\partial}{\partial t_{d_n, \alpha_n}} \langle 1 \rangle_h.$$

Therefore $\langle 1 \rangle_h$ is the crucial quantity to compute. By means of KdV recursion relations

$$\langle \sigma_1(P)P \rangle_h = 2\langle P^4 \rangle_{h-1} + \frac{1}{2} \sum_{h'=0}^h \langle P^2 \rangle_{h'} \langle P^2 \rangle_{h-h'},$$

it is possible [13] to express $\langle 1 \rangle_h$ as a sum of terms of the form $\langle P^{n_1} \rangle_0 \cdots \langle P^{n_j} \rangle_0 / \langle P^3 \rangle_0^{h+j-1}$ for $1 \leq j \leq 3h-3$ with the constraint $\sum_{k=1}^j n_k = 3(j+h-1)$.

The reduction to the punctured sphere arises also in the evaluation of $\text{Vol}_{WP}(\mathcal{M}_{h,n})$. Indeed, at least in some cases, there is a relationship between $\overline{\mathcal{M}}_{h,n}$, $\overline{\mathcal{M}}_{0,n+3h}$ and their volumes¹. The first example is the geometric isomorphism [14] $\overline{\mathcal{M}}_{1,1} \cong \overline{\mathcal{M}}_{0,4}$, and

$$\text{Vol}_{WP}(\mathcal{M}_{1,1}) = 2\text{Vol}_{WP}(\mathcal{M}_{0,4}). \quad (3)$$

To understand this result it is sufficient to recall that the $\wp$ -function enters in the expression of the uniformizing connection of the once punctured torus $\Sigma_{1,1}$ (note that $\wp$ is a solution of the KdV equation)

$$T_{\Sigma_{1,1}} = \frac{1}{2} (\wp(\tau, z) + c(\tau)),$$

where $c(\tau)$ is the accessory parameter for $\Sigma_{1,1}$. Eq.(3) follows from the fact that $T_{\Sigma_{1,1}}$ is strictly related to the uniformizing connection $T_{\Sigma_{0,4}}$ of the Riemann sphere with four punctures since $\wp$ maps $\Sigma_{1,1}$ two-to-one onto the four punctured Riemann sphere. Let us notice that another isomorphism is [15] $\overline{\mathcal{M}}_{2,0} \cong \overline{\mathcal{M}}_{0,6}$.

There is another way to understand why punctured spheres play a crucial role in 2D gravity. The point is to notice that equal size triangulated Riemann surfaces considered in matrix models can be realized in terms of thrice punctured spheres [16]. This aspect is related

¹The space $\mathcal{M}_{h,n}$ is not affine for $h > 2$. Conversely the space $\mathcal{M}_{0,k}$ is finitely covered by the affine space $V^{(k)}$ defined in (6). Thus for $h > 2$ there are not geometrical isomorphisms between $\overline{\mathcal{M}}_{h,n}$ and $\overline{\mathcal{M}}_{0,n+3h}$. However, in principle, nothing excludes the possibility to express $\text{Vol}_{WP}(\mathcal{M}_{h,n})$ in terms of $\text{Vol}_{WP}(\mathcal{M}_{0,n+3h})$.

to arithmetic surfaces theory [16,17]. In this context one should investigate whether this kind of surfaces have some suitable symmetry to define antiholomorphic involution. This question is important in order to investigate Osterwalder-Schrader positivity. This is connected with the problem of defining the adjoint in higher genus. On the sphere it can be done thanks to the natural antinvolution $z \rightarrow \bar{z}^{-1}$. In higher genus this problem has been solved only on a Schottky double where there is a natural antinvolution [18]. Recently Harvey and González Diéz [19] have considered loci of curves which are prime Galois covering of the sphere. In particular they considered the important case of Riemann surfaces admitting non-trivial automorphisms and showed that there is a birational isomorphism between a subset of the moduli space $\mathcal{M}_h$ and $V^{(n)}$ (defined in (6)).

3. The relation between Liouville and uniformization theory of Riemann surfaces arises in considering the Liouville equation

$$\partial_{\bar{z}} \partial_z \varphi_{cl} = \frac{e^{\varphi_{cl}}}{2}, \quad (4)$$

which is uniquely satisfied by the Poincaré metric (i.e. the metric with Gaussian curvature -1). This metric can be written in terms of the inverse of the uniformizing map J_H , that is $e^{\varphi_{cl}} = |J_H^{-1}|^2 / (\text{Im } J_H^{-1})^2$, $J_H : H \rightarrow \Sigma \cong H/\Gamma$ where H is the upper half-plane and Γ a Fuchsian group. Let us introduce the n -punctured sphere $\Sigma = \hat{\mathbf{C}} \setminus \{z_1, \dots, z_n\}$, $\hat{\mathbf{C}} \equiv \mathbf{C} \cup \{\infty\}$. Its moduli space is the space of classes of isomorphic Σ 's, that is

$$\mathcal{M}_{0,n} = \{(z_1, \dots, z_n) \in \hat{\mathbf{C}}^n | z_j \neq z_k \text{ for } j \neq k\} / \text{Symm}(n) \times PSL(2, \mathbf{C}), \quad (5)$$

where $\text{Symm}(n)$ acts by permuting $\{z_1, \dots, z_n\}$ whereas $PSL(2, \mathbf{C})$ acts by linear fractional transformations. By $PSL(2, \mathbf{C})$ we can recover the ‘standard normalization’: $z_{n-2} = 0$, $z_{n-1} = 1$ and $z_n = \infty$. Furthermore, without loss of generality, we assume that $w_{n-2} = 0$, $w_{n-1} = 1$ and $w_n = \infty$. For the classical Liouville tensor we have

$$T^F(z) = \sum_{k=1}^{n-1} \left(\frac{1}{2(z - z_k)^2} + \frac{c_k}{z - z_k} \right), \quad \lim_{z \rightarrow \infty} T^F(z) = \frac{1}{2z^2} + \frac{c_n}{z^3} + \mathcal{O}\left(\frac{1}{|z|^4}\right),$$

with the following constraints on the *accessory parameters*

$$\sum_{k=1}^{n-1} c_k = 0, \quad \sum_{k=1}^{n-1} c_k z_k = 1 - n/2, \quad \sum_{k=1}^{n-1} z_k (1 + c_k z_k) = c_n.$$

The c_k 's are functions on

$$V^{(n)} = \{(z_1, \dots, z_{n-3}) \in \mathbf{C}^{n-3} | z_j \neq 0, 1; z_j \neq z_k, \text{ for } j \neq k\}. \quad (6)$$

Note that

$$\mathcal{M}_{0,n} \cong V^{(n)} / \text{Symm}(n), \quad (7)$$

where the action of $\text{Symm}(n)$ on $V^{(n)}$ is defined by comparing (5) with (7).

Let us now consider the compactification divisor (in the sense of Deligne-Knudsen-Mumford) $D = \overline{V}^{(n)} \setminus V^{(n)}$. This divisor decomposes in the sum of divisors $D_1, \dots, D_{[n/2]-1}$ which are subvarieties of real dimension $2n - 8$. The locus D_k consists of surfaces that split, on removal of the node, into two Riemann spheres with $k + 2$ and $n - k$ punctures. In particular D_k consists of $C(k)$ copies of the space $\overline{V}^{(k+2)} \times \overline{V}^{(n-k)}$ where $C(k) = \binom{n}{k+1}$ for $k = 1, \dots, \frac{(n-1)}{2} - 1$, with the exception that for n even $C(n/2 - 1) = \frac{1}{2} \binom{n}{n/2}$. It turns out that the image of the D_k 's, provide a basis in $H_{2n-8}(\overline{\mathcal{M}}_{0,n}, \mathbf{R})$.

In the case of the punctured Riemann sphere eq.(4) follows from the Liouville action

$$S^{(n)} = \lim_{r \rightarrow 0} \left[\int_{\Sigma_r} (\partial_z \varphi \partial_{\bar{z}} \varphi + e^\varphi) + 2\pi(n \log r + 2(n-2) \log |\log r|) \right],$$

where $\Sigma_r = \Sigma \setminus \left(\bigcup_{i=1}^{n-1} \{z \mid |z - z_i| < r\} \cup \{z \mid |z| > r^{-1}\} \right)$. This action, evaluated on the classical solution, is the Kähler potential for the Weil-Petersson two-form on $V^{(n)}$ [20]

$$\omega_{WP}^{(n)} = \frac{i}{2} \bar{\partial} \partial S_{cl}^{(n)} = -i\pi \sum_{j,k=1}^{n-3} \frac{\partial c_k}{\partial \bar{z}_j} d\bar{z}_j \wedge dz_k. \quad (8)$$

Let us consider the volume of moduli space of punctured Riemann spheres

$$\text{Vol}_{WP}(\mathcal{M}_{0,n}) = \frac{1}{(n-3)!} \int_{\mathcal{M}_{0,n}} \omega_{WP}^{(n) \, n-3} = \frac{1}{(n-3)!} [\omega_{WP}^{(n)}]^{n-3} \cap [\overline{\mathcal{M}}_{0,n}].$$

Recently it has been shown that [21]

$$\text{Vol}_{WP}(\mathcal{M}_{0,n}) = \frac{1}{n!} \text{Vol}_{WP}(V^{(n)}) = \frac{\pi^{2(n-3)} V_n}{n!(n-3)!}, \quad n \geq 4,$$

where $V_n = \pi^{2(3-n)} [\omega_{WP}^{(n)}]^{n-3} \cap [\overline{V}^n]$ satisfies the recursion relations

$$V_3 = 1, \quad V_n = \frac{1}{2} \sum_{k=1}^{n-3} \frac{k(n-k-2)}{n-1} \binom{n}{k+1} \binom{n-4}{k-1} V_{k+2} V_{n-k}, \quad n \geq 4. \quad (9)$$

Remarkably the basic tools in the computation of the volumes are classical Liouville theory and intersection theory.

4. We now consider the differential equation associated with (9). First of all we define

$$a_k = \frac{V_k}{(k-1)((k-3)!)^2}, \quad k \geq 3, \quad (10)$$

so that (9) becomes

$$a_3 = 1/2, \quad a_n = \frac{1}{2} \frac{n(n-2)}{(n-1)(n-3)} \sum_{k=1}^{n-3} a_{k+2} a_{n-k}, \quad n \geq 4. \quad (11)$$

Eq.(11) is equivalent to the differential equation

$$g'' = \frac{g'^2 t - g g' + g' t}{t(t-g)}, \quad (12)$$

where $g(t) = \sum_{k=3}^{\infty} a_k t^{k-1}$. Notice that by (8)

$$g(t) = \sum_{k=3}^{\infty} \frac{k(k-2)}{(k-3)!} t^{k-1} \int_{\mathcal{M}_{0,k}} \left(\frac{i\bar{\partial}\partial S_{cl}^{(k)}}{2\pi^2} \right)^{k-3}, \quad (13)$$

where “ $\int_{\mathcal{M}_{0,3}} 1$ ” $\equiv \frac{1}{6}$. Function $g(t)$ resembles a topological expansion of string theory. Furthermore the structure of eq.(12) resembles the Painlevé I. These remarks indicate that it is possible to recover the specific heat of pure gravity in the continuum. Actually, we will recover the Painlevé I by classical Liouville theory. In particular we will get the recursion relations for the Painlevé I by performing a suitable modification of the Weil-Petersson volume form $\omega_{WP}^{(n) n-3}$. Remarkably, as we will show, it is possible to perform the substitution

$$\omega_{WP}^{(n) n-3} \longrightarrow \omega_{WP}^{(n) n-4} \wedge \omega^F,$$

in (13) without changing the general structure of (11); that is we will get recursion relations of the following structure

$$A_n = C(n) \sum_{k=1}^{n-3} A_{k+2} A_{n-k}, \quad n \geq 4. \quad (14)$$

The first problem is to find a suitable expansion for the Painlevé I field such that the structure of the associated recursion relation be the same of (14). Remarkably this expansion exists, namely

$$f(t) = t^{-12} \sum_{k=3}^{\infty} d_k t^{5k}. \quad (15)$$

It is interesting that in searching the expansion reproducing the general structure of (11), which is a result obtained from continuous Liouville theory, one obtains an expansion involving only *positive* powers of t . With this expansion the Painlevé I

$$f^2(t) - \frac{1}{3} f''(t) = t, \quad (16)$$

is equivalent to the recursion relations²

$$d_n = \frac{3}{(12-5n)(13-5n)} \sum_{k=1}^{n-3} d_{k+2} d_{n-k}, \quad d_3 = -1/2, \quad (17)$$

which has the same structure of (11). We now investigate on the possible volume forms reproducing (17). To understand which kind of modification to $\omega_{WP}^{(n) n-3}$ can be performed without changing the basic structure of (11) we recall basic steps in [21] to obtain (9).

Let D_{WP} be the $(2n-8)$ -cycle dual to the Weil-Petersson class $[\omega_{WP}^{(n)}]$. To compute the volumes it is useful to expand D_{WP} in terms of the divisors D_k in the boundary of moduli space. It turns out that [21]

$$D_{WP} = \frac{\pi^2}{n-1} \sum_{k=1}^{[n/2]-1} k(n-k-2) D_k. \quad (18)$$

Let us set

$$\tilde{V}_n = \pi^{2(n-3)} V_n = [\omega_{WP}^{(n)}]^{n-3} \cap [\overline{V}^{(n)}] = [\omega_{WP}^{(n)}]^{n-4} \cap \left([\omega_{WP}^{(n)}] \cap [\overline{V}^{(n)}] \right).$$

On the other hand $[\omega_{WP}^{(n)}] \cap [\overline{V}^{(n)}] = D_{WP} \cdot \overline{V}^{(n)} = D_{WP}$, so that by (18)

$$\tilde{V}_n = [\omega_{WP}^{(n)}]^{n-4} \cap [D_{WP}] = \frac{\pi^2}{n-1} \sum_{k=1}^{[n/2]-1} k(n-k-2) [\omega_{WP}^{(n)}]^{n-4} \cap [D_k].$$

Since D_k consists of $C(k)$ copies of the space $\overline{V}^{(k+2)} \times \overline{V}^{(n-k)}$, we have

$$\tilde{V}_n = \frac{\pi^2}{n-1} \sum_{k=1}^{[n/2]-1} k(n-k-2) C(k) [\omega_{WP}^{(n)}]^{n-4} \cap [\overline{V}^{(k+2)} \times \overline{V}^{(n-k)}].$$

Finally, since [21]

$$[\omega_{WP}^{(n)}]^{n-4} \cap [\overline{V}^{(k+2)} \times \overline{V}^{(n-k)}] = [\omega_{WP}^{(k+2)} + \omega_{WP}^{(n-k)}]^{n-4} \cap [\overline{V}^{(k+2)} \times \overline{V}^{(n-k)}], \quad (19)$$

it follows that

$$\tilde{V}_3 = 1, \quad \tilde{V}_n = \frac{\pi^2}{n-1} \sum_{k=1}^{[n/2]-1} k(n-k-2) C(k) \binom{n-4}{k-1} \tilde{V}_{k+2} \tilde{V}_{n-k}, \quad n \geq 4,$$

which coincides with (9).

²Notice that $(-1)^k d_k$ is positive.

We now introduce the divisor

$$D^F = \frac{\pi^2}{n-1} \sum_{k=1}^{[n/2]-1} k(n-k-2) F(n, k) D_k, \quad (20)$$

where $F(n, k)$ is a function to be determined. Let $[\omega^F]$ be the Poincaré dual to D^F and define

$$Z_n^F = \int_{\overline{\mathcal{M}}_{0,n}} \omega_{WP}^{(n)}{}^{n-4} \wedge \omega^F = \int_{\overline{\mathcal{M}}_{0,n}} \left(\frac{i\bar{\partial}\partial S_{cl}^{(n)}}{2} \right)^{n-4} \wedge \omega^F, \quad n \geq 4. \quad (21)$$

An important aspect of (21) is that we can use the recursion relations (11) to obtain non-perturbative results. This possibility is based on the obvious, but important fact, that $[\omega_{WP}^{(n)}]^{n-3} \cap [\overline{V}^{(n)}] = [\omega_{WP}^{(n)}]^{n-4} \cap [D_{WP}]$ implying that the general structure of (11) (the same of (17)) is unchanged under the substitution $\omega_{WP}^{(n)}{}^{n-3} \longrightarrow \omega_{WP}^{(n)}{}^{n-4} \wedge \omega^F$. To see this note that

$$Z_n^F = \frac{1}{n!} [\omega_{WP}^{(n)}]^{n-4} \cap [D^F] = \frac{\pi^2}{(n-1)n!} \sum_{k=1}^{[n/2]-1} F(n, k) k(n-k-2) [\omega_{WP}^{(n)}]^{n-4} \cap [D_k], \quad (22)$$

On the other hand by (19)

$$\sum_{k=1}^{[n/2]-1} F(n, k) k(n-k-2) [\omega_{WP}^{(n)}]^{n-4} \cap [D_k] = \frac{1}{2} \sum_{k=1}^{n-3} F(n, k) \frac{k(n-k-2)}{n-1} \binom{n}{k+1} \binom{n-4}{k-1} V_{k+2} V_{n-k}, \quad (23)$$

and by (10)

$$Z_n^F = \frac{\pi^2}{2} \frac{(n-4)!}{n-1} \sum_{k=1}^{n-3} F(n, k) a_{k+2} a_{n-k}, \quad n \geq 4. \quad (24)$$

Let us define the ‘Liouville F -models’

$$\mathcal{Z}^{F,\alpha}(x) = x^{-\alpha} \sum_{k=3}^{\infty} x^k Z_k^F, \quad (25)$$

where x is the coupling constant. These models are classified by α , $F(n, k)$ and Z_3^F . We now show that $\mathcal{Z}^{F,\alpha}(x)$ includes pure gravity. In fact, putting

$$\mathcal{Z}(t) = \mathcal{Z}^{F_0, \alpha_0}(t^5), \quad Z_3^{F_0} = -1/2, \quad \alpha_0 = 12/5, \quad (26)$$

where

$$F_0(n, k) = \frac{6}{\pi^2} \frac{(n-1)}{(12-5n)(13-5n)(n-4)!} \frac{Z_{k+2}^{F_0} Z_{n-k}^{F_0}}{a_{k+2} a_{n-k}}, \quad (27)$$

we have, by (24) and (27),

$$Z_3^{F_0} = -1/2, \quad Z_n^{F_0} = \frac{3}{(12-5n)(13-5n)} \sum_{k=1}^{n-3} Z_{k+2}^{F_0} Z_{n-k}^{F_0}, \quad n \geq 4, \quad (28)$$

so that by (15)-(17)

$$\mathcal{Z}(t) = t^{-12} \sum_{k=4}^{\infty} t^{5k} \int_{\mathcal{M}_{0,k}} \left(\frac{i\bar{\partial}\partial S_{cl}^{(k)}}{2} \right)^{k-4} \wedge \omega^F - \frac{t^3}{2}, \quad (29)$$

satisfies the Painlevé I

$$\mathcal{Z}^2(t) - \frac{1}{3}\mathcal{Z}''(t) = t. \quad (30)$$

5. In conclusion we have introduced a class of Liouville models by defining a suitable D^F divisor. These Liouville F -models (LFM) include pure gravity. In this context we recall that the Liouville action arises also in the correlators of topological gravity [8].

Punctures correspond to real points on the boundary of the upper half-plane. Correspondingly one can define hyperelliptic Riemann surfaces. In the case of infinite genus one gets the McKean-Trubowitz [22] model which is related to matrix model. This suggests a nonperturbative formulation on H with the image of punctures related to the eigenvalues of the Hermitian matrix models. In the discrete version of this approach one should be able to connect this formulation with the ideas at the basis of [23].

Acknowledgements I would like to thank G. Bonelli for stimulating discussions.

References

- [1] E. Brézin and V. Kazakov, Phys. Lett. **236B** (1990) 144. M. Douglas and S. Shenker, Nucl. Phys. **B335** (1990) 635. D. Gross and A. Migdal, Phys. Rev. Lett. **64** (1990) 127.
- [2] L. Alvarez-Gaumé, *Random Surfaces, Statistical Mechanics And String Theory*, Lausanne lectures, winter 1990. P. Ginsparg, *Matrix Models Of 2D Gravity*, Trieste Lectures, LA-UR-91-4101, hep-th/9112013. E. Martinec, *An Introduction To 2D Gravity And Solvable String Models*, Trieste Lectures, RU-91-51, hep-th/9112019. A. Morozov, *Integrability And Matrix Models*, ITEP-M2/93, hep-th/9303139. P. Di Francesco, P. Ginsparg and J. Zinn-Justin, *2D Gravity And Random Matrices*, LA-UR-93-1722, SPhT/93-061, hep-th/9306153.

- [3] A.M. Polyakov, Phys. Lett. **103B** (1981) 207.
- [4] F. David, Mod. Phys. Lett **A3** (1988) 509. J. Distler and H. Kawai, Nucl. Phys. **B321** (1989) 509.
- [5] J.-L. Gervais, Comm. Math. Phys. **130** (1990) 257; **138** (1991) 301.
- [6] M. Matone, *Quantum Riemann Surfaces, 2D Gravity And The Geometrical Origin Of Minimal Models*, preprint DFPD/93/TH/62, hep-th/9309096.
- [7] L.A. Takhtajan, *Liouville Theory: Quantum Geometry Of Riemann Surfaces*, hep-th/9308125.
- [8] M. Matone, *Uniformization Theory and 2D Gravity. I. Liouville Action And Intersection Numbers*, IC-MATH/8-92, DFPD-TH/92/41, hep-th/9306150.
- [9] E. Witten, Nucl. Phys. **B340** (1990) 281; Surv. Diff. Geom. **1** (1991) 243. R. Dijkgraaf and E. Witten, Nucl. Phys. **B342** (1990) 486.
- [10] V.G. Knizhnik, Sov. Phys. Usp. **32** (11) (1989) 945.
- [11] J.J. Duistermaat and G.J. Heckman, Invent. Math. **69** (1982) 259; **72** (1983) 153. N. Berline and M. Vergne, Duke Math. J. **50** (1983) 539. R.F. Picken J. Math. Phys. **31** (1990) 616.
- [12] D. Friedan and S. Shenker, Nucl. Phys. **B281** (1987) 509.
- [13] J.H. Horne and S.P. Martin, Phys. Lett. **258B** (1991) 322. A. Sen, Int. J. Mod. Phys. **A7** (1992) 2559. K. Li, Nucl. Phys. **B354** (1991) 725.
- [14] S.A. Wolpert, Ann. of Math. **118** (1983) 491.
- [15] J. Igusa, Ann. of Math. **72** (1960) 612.
- [16] A. Levin and A. Morozov, Phys. Lett. **243B** (1990) 207.
- [17] D.-J. Smit, Comm. Math. Phys. **143** (1992) 253.
- [18] A. Jaffe, S. Klimek and A. Lesniewski, Comm. Math. Phys. **126** (1989) 421.

- [19] G. González-Diéz and W.J. Harvey, *Moduli Of Riemann Surfaces With Symmetry*, in Discrete Groups and Geometry, ed. W.J. Harvey and C. Maclachlan, Cambridge 1992. G. González-Diéz Proc. London Math. Soc. 62 (1991) 469; *On Prime Galois Covering Of The Riemann Sphere*, preprint.
- [20] P.G. Zograf and L.A. Takhtajan, Math. USSR Sbornik, **60** (1988) 143; **60** (1988) 297. L.A. Takhtajan, Proc. Symp. Pure Math. (AMS) **49**, part 1 (1989) 581.
- [21] P.G. Zograf, *The Weil-Petersson Volume Of The Moduli Space Of Punctured Spheres*, to appear in Cont. Math..
- [22] H.P. McKean and E. Trubowitz, Comm. Pure Appl. Math. **29** (1976) 143.
- [23] L. Bonora and C.S. Xiong, *Correlation Functions Of Two-Matrix Models*, SISSA 172/93/EP, BONN-HE-45/93, hep-th/9311089.